

## **SUP-ICU trial**

## **Inclusion criteria**

- ✓ Acute (unplanned) admission to the ICU
- ✓ Aged 18 years or above
- ✓ One or more of the following risk factors:
  - Shock (continous infusion with vasopressor or inotrope, mean arterial blood pressure < 70 mmHg, systolic blood pressure < 90 mmHg or lactate > 4 mmol/l)
  - Acute or chronic intermittent or continuous renal replacement therapy
  - Invasive mechanical ventilation which is expected to last > 24 hours. When in doubt of the forecast, the patient should be enrolled
  - Coagulopathy (platelets < 50 x 10<sup>9</sup>/l, INR > 1.5 or PT > 20 seconds)
  - Ongoing treatment (not prophylaxis) with anticoagulant medication
  - History of coagulopathy (platelets < 50 x 109/l or INR > 1.5 or PT > 20 seconds within 6
    months of this hospital admission
  - History of chronic liver disease (history of portal hypertension, variceal bleeding or hepatic encephalopathy or cirrhosis proven by biopsy, CT scan or ultrasound)

## **Exclusion criteria**

- Contraindications to PPI (including intolerance of PPI and treatment with atazanavir (HIV medication))
- Current <u>daily</u> treatment with PPI and/or H2RA
- **✗** GI bleeding of any origin during current hospital admission
- Diagnosed with peptic ulcer during current hospital admission
- Organ transplant during current hospital admission
- Withdrawal from active therapy or brain death
- Fertile woman with positive urine human chorionic gonadotropin (hCG) or plasma-hCG
- **×** Consent according to national regulations not obtainable