| Baseli | ne (0/28) | |
|----------|--|---|
| Title: I | Baseline | |
| Instruc | ions: | |
| | | BASELINE |
| | | |
| | | Hotline: +45 35 45 06 06 |
| | | General patient information |
| | Ra | indomisation timestamp: |
| | | |
| BL1 | Sex? | O Male O Female [info] |
| BL2 | Hospital admission date? | (dd-mm-yyyy) [info] |
| BL3 | ICU admission date? | (dd-mm-yyyy) [info] |
| BL4 | ICU admission time? | (24 hours, hh:mm) [info] |
| BL5 | From where was the patient admitted? | Emergency department or directly from the pre-hospital setting [info] Hospital ward Operating or recovery room Another ICU |
| BL6 | Focus of infection (documented or suspected): | Pulmonary [info] Gastrointestinal Urinary tract Skin or soft tissue Other |
| | Co-m | orbidities prior to ICU admission |
| BL7 | History of ischemic heart disease or heart failure? | ○ Yes ○ No [info] |
| BL8 | History of chronic hypertension? | ○ Yes ○ No [info] |
| BL9 | Chronic renal replacement therapy? | ○ Yes ○ No [info] |
| | | es, interventions and vital parameters Indomisation timestamp: |
| BL10 | Participant weight | (kg) [info] |
| BL11 | Highest plasma lactate value in the last 3 hours prior to randomisation | mmol/L mg/dL [info] (mmol/L) (mg/dL) |
| BL12 | Highest dose of noradrenaline in the last 3 hours prior to randomisation: | (µg/kg/min) [info] |
| BL13 | Infused volume of IV fluids in the last 24 hours prior to randomisation: | (mL) [info] |
| BL14 | Use of systemic (IV, IM or oral/per GI tube) corticosteroids within 24 hours prior to randomisation: | ○ Yes ○ No [info] |
| BL15 | Highest plasma creatinine value within the last 24 hours prior to randomisation: | ○ µmol/L ○ mg/dL □ Not Available (µmol/L) (mg/dL) |

| BL16 | Use of <u>acute</u> renal replacement therapy in the last <u>3 days</u> prior to randomisation: | ○ Yes ○ No [info] |
|--------------|---|---|
| BL17 | Use of <u>any form</u> of renal replacement therapy within the <u>last 24</u> hours prior to randomisation: | ○ Yes ○ No [info] |
| BL18 | Habitual plasma creatinine value prior to current hospitalisation: estimated or measured. | μmol/L |
| | | Measured Estimated |
| | | ed Mortality Score for the Intensive Care Unit) |
| | | |
| BL19 | | ed Mortality Score for the Intensive Care Unit) |
| BL19 BL20 | Lowest systolic blood pressure within the last 24 hours prior to | ed Mortality Score for the Intensive Care Unit) andomisation timestamp: |